



**QUEEN'S
UNIVERSITY
BELFAST**

Quacks snack on smacks: mallard ducks (*Anas platyrhynchos*) observed feeding on hydrozoans (*Velella velella*)

Phillips, N., Eagling, L., Harrod, C., Cappanera, V., Houghton, J., & Reid, N. (2017). Quacks snack on smacks: mallard ducks (*Anas platyrhynchos*) observed feeding on hydrozoans (*Velella velella*). *Plankton & Benthos Research*, 12(2), 143-144. https://www.jstage.jst.go.jp/article/pbr/12/2/12_P120207/_article

Published in:
Plankton & Benthos Research

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2017 Plankton Society of Japan.
This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Quacks snack on smacks: mallard ducks (*Anas platyrhynchos*) observed feeding on hydrozoans (*Velella velella*)

Natasha Phillips^{*a}, Lawrence Eagling^b, Chris Harrod^c, Neil Reid^{a,d}, Valentina Cappanera^e, Jonathan Houghton^{*a,b,c}

^a*School of Biological Sciences, Queen's University Belfast MBC Building, 97 Lisburn Road, Belfast, BT9 7BL, U.K.*, ^b*Queen's Marine Laboratory, 12-13 The Strand Portaferry, Newtownards BT22 1PF*, ^c*Instituto de Ciencias Naturales Alexander von Humboldt, Universidad de Antofagasta, Avenida Angamos 601, Antofagasta, Chile*, ^d*Institute of Global Food Security, Queen's University Belfast, 18-30 Malone Road, Belfast, BT9 5BN*, ^e*Area Marina Protetta di Portofino, Ministry for the Environment and for the Protection of Territory and Sea, Santa Margherita Ligure, Genova, Italy*

*Author to whom correspondence should be addressed: Natasha Phillips, Queen's University Belfast, Medical Biology Centre, 97 Lisburn Road, Belfast, BT9 7AE, nphillips01@qub.ac.uk

Running page head

Mallard ducks observed feeding on hydrozoans

Abstract

This study presents new evidence of the extensive trophic role of gelatinous zooplankton by documenting typically non-marine predators, mallard ducks, feeding on hydrozoans in shallow, coastal environments.

Keywords

Avian; cryptic trophic linkages; gelatinous zooplankton; predator-prey; trophic pathways; *Velella velella*

Far from being trophic dead ends, gelatinous zooplankton are now known to play a host of diverse roles in ecosystem functioning (Doyle et al. 2014); from mass “jelly falls” following jellyfish blooms that act as carbon sinks (Sweetman & Chapman 2015), through to the provision of shelter for developing invertebrate and fish communities (e.g. Lynam & Brierley 2007, D’Ambra et al. 2014, Fleming et al. 2014). Moreover, aside from specialised obligate gelativores such as leatherback turtles (*Dermochelys coriacea* Vandelli, 1761; Brongersma 1969), growing evidence suggests that a vast array of taxa also consume gelatinous prey periodically (see reviews: Arai 2005, Ates 2017). Predators of note include juvenile bluefin tuna (*Thunnus thynnus* Linnaeus, 1758; Cardona et al. 2012), Atlantic bumper (*Chloroscombrus chrysurus* Linnaeus, 1766; D’Ambra et al. 2014), spiny lobsters (*Panulirus interruptus* Randall, 1840; O’Rorke et al. 2014), deep-sea 7-armed octopus (*Haliphron atlanticus* Steenstrup, 1861; Hoving & Haddock 2017) and other gelatinous species (e.g. Purcell 1981, Purcell 1991).

By contrast, episodic feeding on gelatinous species by avian predators has only been described recently, with initial reports of scyphozoan jellyfish acting as fish aggregation devices (e.g. Richardson et al. 2009, Sato et al. 2015) which can be exploited by birds. With the advent of animal-borne cameras there is also evidence of penguins *Pygoscelis adeliae* (Hombron & Jacquinot, 1841) feeding directly on gelatinous prey throughout the Southern Ocean (Thiebot et al. 2016). Here we build on this growing body of evidence by providing observational evidence of an unreported trophic pathway; the ingestion of the hydrozoan *Velella velella* (Linnaeus, 1758) by mallard ducks *Anas platyrhynchos* (Linnaeus, 1758).

Following a period of unsettled weather in the Italian district of Liguria in late May 2016, two individuals (one male, one female) were observed browsing and feeding on a large

patch of *V. velella* (Fig 1) which had been washed into the shallow harbour of Santa Margherita Ligure (44°20'1.1"N, 9°12'50.7"E on 30th May 2016, see Fig 2).

The ducks were seen feeding on *V. velella* within the hydroid patch (Fig 3); but poor lighting at the time prevented any estimates of ingestion rate or prey handling duration. Since we are not proposing that ducks feed routinely on such prey, the simple finding that such trophic links even exist is not hindered by a lack of empirical data. More explicitly, although predation on *V. velella* has been noted in a variety of oceanic vertebrate predators (Purcell et al. 2012) including several other avian species such as fulmars (*Fulmarus glacialis* Linnaeus, 1761; Williams et al. 1991), predation by typically non-marine species illustrates that the trophic role of gelatinous zooplankton can, on occasion, extend further than previously thought.

Acknowledgements

The authors would like to acknowledge the generous contribution of all grant providers and funding bodies who supported this project; The Alice McCosh Trust, Marine Institute, Society of Experimental Biology, The Sir Thomas Dixon Travel Grant, The MacQuitty Grant Scholarship and The Royal Society of Biology. N.D.P was funded by The Fisheries Society of the British Isles. All applicable international, national, and institutional guidelines for the care and use of animals were followed. This article does not contain any studies with human participants performed by any of the authors. The authors declare that they have no conflict of interest.

References

- Arai M (2005) Predation on pelagic coelenterates: a review. J Mar Biol Ass UK 85:523-536.
- Ates RML (2017) Benthic scavengers and predators of jellyfish, material for a review. Plank Benth Res 12:71-77.

76 Brongersma LD (1969) Miscellaneous notes on turtles. Proc K Ned Akad Wet C 72:90-102.

77 Cardona L, Álvarez de Quevedo I, Borrell A, Aguilar A (2012) Massive consumption of gelatinous plankton
78 by Mediterranean apex predators PLoS ONE 7: e31329.

79 D'Ambra I, Graham WM, Carmichael RH, Hernandez FJ Jr. (2015) Fish rely on scyphozoan hosts as a
80 primary food source: evidence from stable isotope analysis. Mar Biol 162:247-252.

81 Doyle TK, Hays GC, Harrod C, Houghton JDR (2014) Ecological and societal benefits of jellyfish. In: Pitt
82 KA and Lucas CH (eds) Jellyfish Blooms. Springer Science and Business Media Dordrecht, 105-122.

83 ESRI 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.

84 Fleming NE, Harrod C, Griffin D, Newton J, Houghton JDR (2014) Scyphozoan jellyfish provide short-term
85 reproductive habitat for hyperiid amphipods in a temperate near-shore environment. Mar Ecol Prog Ser
86 510: 229-240.

87 Hoving HJT, Haddock SHD (2017) The giant deep-sea octopus *Haliphron atlanticus* forages on gelatinous
88 fauna. Sci Rep 7:44952.

89 Lynam CP, Brierley AS (2007) Enhanced survival of 0-group gadoid fish under jellyfish umbrellas. Mar Biol
90 150:1397-1401.

91 Purcell JE, Clarkin E and Doyle TK (2012) Foods of *Velella velella* (Cnidaria: Hydrozoa) in algal rafts and
92 its distribution in Irish seas. Hydrobiologia 690:47-55.

93 Purcell JE (1981) Dietary composition and diel feeding patterns of epipelagic siphonophores. Mar Biol
94 65:83-90.

95 Purcell JE (1991) A review of cnidarians and ctenophores feeding on competitors in the plankton.
96 Hydrobiologia 216:335-342.

97 O'Rourke R, Lavery SD, Wang M, Gallego R, Waite AM, Beckley LE, Thompson PA, Jeffs AG (2014)
98 Phyllosoma associated with large gelatinous zooplankton: hitching rides and stealing bites. ICES J Mar
99 Sci 72:124-127.

100 Richardson AJ, Bakun A, Hays GC, Gibbons MJ (2009) The jellyfish joyride: causes, consequences and
101 management responses to a more gelatinous future. Trends Ecol Evol 24:312-322.

102 Sato NN, Kokubun N, Yamamoto T, Watanuki Y, Kitaysky AS, Takahashi A (2015) The jellyfish buffet:
103 jellyfish enhance seabird foraging opportunities by concentrating prey. Biol Lett 11:20150358.

Sweetman AK, Chapman A (2015) First assessment of flux rates of jellyfish carcasses (jelly-falls) to the benthos reveals the importance of gelatinous material for biological C-cycling in jellyfish-dominated ecosystems. Front Mar Sci 2:47.

Thiebot JB, Ito K, Raclot T, Poupart T, Kato A, Ropert-Coudert Y, Takahasi A (2016) On the significance of Antarctic jellyfish as food for Adélie penguins, as revealed by video loggers. Mar Biol 163:108.

Williams RB, Cornelius PFS, Hughes RG, Robson EA (1991) Coelenterate Biology; Recent research on cnidarian and ctenophore. Hydrobiologia 305: 216-217.

Figure legends

Fig. 1 A by-the-wind-sailor or *Velella velella* washed up on Santa Margherita Ligure beach



Fig. 2 Map showing location of Santa Margherita Ligure, Italy produced using ARCGIS 10.3.1 (ESRI, California, USA)




Fig. 3 Mallard ducks feeding on *Velella velella* in Santa Margherita Ligure harbour



121